

Manufacturer/S	upplier:		
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD			
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District			
Wuxi Jiangsu Province, China			
TELE: 86-510-82732223 FAX: 86-510-82720101-8014			
Web : <u>www.dldevelop.com</u>			
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>			
Product name:	Anti-Human CD57 Monoclonal Antibody(AF647		
	Conjugated)		
Catalog:	DL20837F		
Synonyms:	Galactosylgalactosylxylosylprotein 3-beta-glucuronosyltransferase 1,B3GAT1,GlcAT-P,GlcUAT-P,GLCATP		
Background: CD57, also known as HNK-1, NK-1, and Leu-7 is			
100-115 kD oligosaccharide antigenic determinant			
	expressed on a variety of proteins, lipids, and		
chondroitin sulfate proteoglycans. CD57 is expressed			
on a subset of peripheral blood lymphocytes,			
	including NK cells and CD8+ T cells, and is also expressed on neural cells and striated muscle. CD57 is not expressed on red blood cells, granulocytes,		
	monocytes, or platelets. While the function of CD57		



	is unknown, binding to L-selectin, P-selectin, and a		
	fragment of laminin suggests that CD57 may be involved in cell-matrix interactions. CD57 is increased in some disease states associated with CD4/CD8 imbalances (AIDS, autoimmune disease, viral infections, and allograft transplants).		
Form: Liquid		Isotype: Mouse IgM	
Size: 20 Tests/50	Tests/100	Host: Mouse	
Tests/200 Tests			
Reactivity: Huma	n	Application: FCM	
Concentration:		Conjugation: AF647	
SwissProt:	Q9P2W7		
Buffer:	PBS with 0.05% Proclin300, 1% BSA		

Recommended Use:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the



appropriate dilutions must be determined for individual use.

Data:

Human peripheral blood lymphocytes are stained with Anrti-Human CD57 Monoclonal Antibody(AF647 Conjuaged)(filled gray histogram). Unstained lymphocytes (empty black histogram) are used as control.

Storage:

Keep as concentrated solution.

Store at $2 \sim 8^{\circ}$ C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.

