

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Purified Anti-Human CD58 Monoclonal Antibody

DL20838F **Catalog:**

Synonyms: Lymphocyte function-associated antigen

3,CD58,Ag3,Surface glycoprotein LFA-3,LFA3

Background: CD58, also known as lymphocyte

function-associated antigen 3 (LFA-3) is a 45-70 kD

cell surface protein that is a member of the

immunoglobulin superfamily. Alternative splicing of

CD58 gives rise to transmembrane and

glycosylphosphatidylinositol (GPI)-anchored forms

on cell surface. CD58 is expressed on both

hematopoietic and non-hematopoietic cells including B cells, T cells, monocytes, erythrocytes, endothelial cells, epithelial cells, and fibroblasts. High levels are observed on memory T cells and dendritic cells.



CD58 expressed on antigen presenting cells and target cells enhances T cell recognition via the binding of it's cognate ligand, CD2, on the T cell surface.

Form: Liquid **Isotype:** Mouse IgG1, κ

Size: 25µg/100µg Host: Mouse

Reactivity: Human **Application:** FCM,IP

Concentration: 0.5 mg/mL Conjugation: Unconjugated

SwissProt: P19256

Buffer: Phosphate buffered solution, pH 7.2, containing

0.09% stabilizer.

Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is $\leq 1.0 \,\mu g$ per 10^6 cells in $100 \,\mu L$ volume or $100 \,\mu L$ of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.



Storage:

Keep as concentrated solution.

Store at 2~8°C. Do not freeze.

This product is guaranteed up to one year from purchase.