

Manufacturer/Supplier:			
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD			
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District			
Wuxi Jiangsu Province, China			
TELE: 86-510-82732223 FAX: 86-510-82720101-8014			
Web : <u>www.dldevelop.com</u>			
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>			
Product name:	Anti-Human CD58 Monoclonal Antibody(APC		
	Conjugated)		
Catalog:	DL20841F		
Synonyms:	Lymphocyte function-associated antigen		
	3,CD58,Ag3,Surface glycoprotein LFA-3,LFA3		
<b>Background:</b>	CD58, also known as lymphocyte		
	function-associated antigen 3 (LFA-3) is a 45-70 kD		
	cell surface protein that is a member of the		
	immunoglobulin superfamily. Alternative splicing of		
	CD58 gives rise to transmembrane and		
	glycosylphosphatidylinositol (GPI)-anchored forms		
	on cell surface. CD58 is expressed on both		
	hematopoietic and non-hematopoietic cells including		
	B cells, T cells, monocytes, erythrocytes, endothelial		
cells, epithelial cells, and fibroblasts. High levels are			



observed on memory T cells and dendritic cells. CD58 expressed on antigen presenting cells and target cells enhances T cell recognition via the binding of it's cognate ligand, CD2, on the T cell surface.

Form: Liquid		<b>Isotype:</b> Mouse IgG1, κ
Size: 20Tests/100Tests/100Tests×2 Reactivity: Human		Host: Mouse
		Application: FCM
Concentration: 5 µL		Conjugation: APC
SwissProt:	P19256	
Duffor	Dhosphata huffar	ad solution nH 7.2 contain

Buffer:Phosphate buffered solution, pH 7.2, containing<br/>0.09% stabilizer and 1% protein protectant.

## **Recommended Use:**


Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined

## **DEVELOP**

for individual use.

## Data:

Human peripheral blood lymphocytes are stained with Anti-Human CD58 Monoclonal Antibody(APC Conjugated)(filled gray histogram). Unstained lymphocytes (empty black histogram) are used as control.



## Storage:

Keep as concentrated solution.

Store at  $2\sim 8^{\circ}$ C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.