

**Manufacturer/Supplier:**

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

**Address:** A1-203 Mingpin CityII, No.8 Xihudong Road, Liangxi District  
Wuxi Jiangsu Province, China

TELE: 86-510-82732223

FAX: 86-510-82720101-8014

**Web :** [www.dldevelop.com](http://www.dldevelop.com)

Email: [info@dldevelop.com.cn](mailto:info@dldevelop.com.cn)    [service@dldevelop.com.cn](mailto:service@dldevelop.com.cn)

**Product name:** Purified Anti-Human CD74 Monoclonal Antibody

**Catalog:** DL20860F

**Synonyms:** HLA class II histocompatibility antigen gamma chain, CD74, Ia antigen-associated invariant chain, Ii, p33, DHLAG

**Background:** CD74 is a type II transmembrane glycoprotein also known as MHC class II associated invariant chain, invariant chain, Ii, MHC class II chaperone, and MIF receptor. CD74 exists in four isoforms with molecular masses of 33, 35, 41, and 43 kD, depending on genetic splicing. CD74 is primarily expressed on antigen presenting cells, including B cells, monocytes/macrophages, dendritic cells, and Langerhans cells. It is also expressed by activated T cells and activated endothelial and epithelial cells as

well as carcinomas of lung, renal, gastric and thymic origin. The primary function of CD74 is intracellular sorting of MHC class II molecules and regulation of exogenous peptide loading onto MHC class II. It is also involved in the modulation of B cell differentiation and positive selection of CD4+ T cells. It has been reported that CD74 binds MIF (macrophage migration inhibitory factor) and signals through CD44 to regulate innate and adaptive immunity. It is also reported that H. pylori infection occurs through urease B binding of CD74 on gastric epithelial cells, inducing gastric epithelial cell apoptosis, NF- $\kappa$ B activation, and IL-8 production.

**Form:** Liquid

**Isotype:** Mouse IgG1,  $\kappa$

**Size:** 25 $\mu$ g/100 $\mu$ g

**Host:** Mouse

**Reactivity:** Human

**Application:** FCM,IHC,IF,IP

**Concentration:** 0.5 mg/mL

**Conjugation:** Unconjugated

**SwissProt:** P04233

**Buffer:** Phosphate buffered solution, pH 7.2, containing

0.09% stabilizer.

### **Recommended Use:**

Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is  $\leq 2.0 \mu\text{g}$  per  $10^6$  cells in  $100 \mu\text{L}$  volume or  $100 \mu\text{L}$  of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

### **Storage:**

Keep as concentrated solution.

Store at  $2\sim 8^\circ\text{C}$ . Do not freeze.

This product is guaranteed up to one year from purchase.