

| Manufacturer/Supplier:   |   |  |
|--|---|--|
| WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD                         |   |  |
| Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District |   |  |
| Wuxi Jiangsu Province, China                                       |   |  |
| TELE:       86-510-82732223       FAX:       86-510-82720101-8014  |   |  |
| Web : <u>www.dldevelop.com</u>                                     |   |  |
| Email: info@dldevelop.com.cn service@dldevelop.com.cn              |   |  |
|  |   |  |
| Product name:  | Anti-Human CD81 Monoclonal                          |  |
|  | Antibody(PE/Cyanine5 Conjugated)                    |  |
| Catalog:   | DL20876F  |  |
|  |   |  |
| Synonyms:  | CD81 antigen,CD81,26 kDa cell surface protein       |  |
|  | TAPA-1, Target of the antiproliferative antibody    |  |
|  | 1.Tetraspanin-28,Tspan-28,APA1, TSPAN28             |  |
| Background:  | CD81 is a 26 kD non-glycosylated member of the      |  |
|  | tetraspanin superfamily (TM4SF), also known as      |  |
|  | TAPA-1 (target of an antiproliferative antibody).   |  |
|  | CD81 is expressed on T and B cells, NK cells,       |  |
|  | monocytes, dendritic cells, thymocytes, endothelial |  |
|  | cells, and fibroblasts. It also has low levels of   |  |
|  | expression on granulocytes. CD81 induces B cell     |  |
|  | adhesion via VLA-4 integrin and has been shown to   |  |
|  | play a role in early T cell development. CD81       |  |
|  |   |  |



associates with several other cell- surface proteins in a multimolecular complex, including CD19, CD21, CD20, CD37, CD53, and CD82 in B cells, and CD4, CD8 and CD82 in T cells.

| <b>Isotype:</b> Mouse IgG1, κ |
|-------------------------------|
| Host: Mouse Application: FCM  |
|                               |
|                               |
|                               |

**Buffer:** PBS with 0.05% Proclin300, 1% BSA

## **Recommended Use:**

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ l staining volume or per 100  $\mu$ l of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

## **DEVELOP**.

## **Storage:**

Keep as concentrated solution.

Store at  $2\sim 8^{\circ}$ C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.

