

| Manufacturer/Supplier: | | | |
|---|---|--|--|
| WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD | | | |
| Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District | | | |
| Wuxi Jiangsu Province, China | | | |
| TELE: 86-510-82732223 FAX: 86-510-82720101-8014 | | | |
| Web : <u>www.dldevelop.com</u> | | | |
| Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u> | | | |
| | | | |
| Product name: | Anti-Human CD81 Monoclonal Antibody(AF488 | | |
| | Conjugated) | | |
| Catalog: | DL20880F | | |
| | | | |
| Synonyms: | CD81 antigen,CD81,26 kDa cell surface protein TAPA-1,Target of the antiproliferative antibody 1.Tetraspanin-28,Tspan-28,APA1, TSPAN28 | | |
| | | | |
| | | | |
| | | | |
| Background: | CD81 is a 26 kD non-glycosylated member of the | | |
| | tetraspanin superfamily (TM4SF), also known as | | |
| | TAPA-1 (target of an antiproliferative antibody). | | |
| | CD81 is expressed on T and B cells, NK cells, | | |
| | monocytes, dendritic cells, thymocytes, endothelial | | |
| | cells, and fibroblasts. It also has low levels of | | |
| | expression on granulocytes. CD81 induces B cell | | |
| | adhesion via VLA-4 integrin and has been shown to | | |
| | play a role in early T cell development. CD81 | | |
| | | | |



associates with several other cell- surface proteins in a multimolecular complex, including CD19, CD21, CD20, CD37, CD53, and CD82 in B cells, and CD4, CD8 and CD82 in T cells.

| Form: Liquid | | Isotype: Mouse IgG1, κ |
|---|---|-------------------------------|
| Size: 20Tests/100Tests/100Tests×2 Reactivity: Human | | Host: Mouse |
| | | Application: FCM |
| Concentration: 5 µL | | Conjugation: AF488 |
| SwissProt: | P60033 | |
| Buffer: | Phosphate buffered solution, pH 7.2, containing | |

Recommended Use:


Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

0.09% stabilizer and 1% protein protectant.

DEVELOP

Data:

Human peripheral blood lymphocytes are stained with Anti-Human CD81 Monoclonal Antibody (AF488 Conjugated)(filled gray histogram). Unstained lymphocytes (empty black histogram) are used as control.



Storage:

Keep as concentrated solution.

Store at $2\sim 8^{\circ}$ C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.