

**Manufacturer/Supplier:**

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

**Address:** A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District  
Wuxi Jiangsu Province, China

TELE: 86-510-82732223

FAX: 86-510-82720101-8014

**Web :** [www.dldevelop.com](http://www.dldevelop.com)

Email: [info@dldevelop.com.cn](mailto:info@dldevelop.com.cn)    [service@dldevelop.com.cn](mailto:service@dldevelop.com.cn)

**Product name:** Purified Anti-Human CD13 Monoclonal Antibody

**Catalog:** DL20882F

**Synonyms:** Aminopeptidase N, ANPEP, AP-N, hAPN, Alanyl aminopeptidase, Aminopeptidase M, AP-M, Microsomal aminopeptidase, gp150, CD13

**Background:** CD13 is a 150-170 kD type II transmembrane glycoprotein also known as aminopeptidase N, APN, and gp150. This zinc metallopeptidase is expressed as a homodimer on granulocytes, myeloid progenitors, endothelial cells, epithelial cells and subset of granular lymphoid cells. It is not expressed on platelets or erythrocytes. CD13 is thought to be involved in the metabolism of many regulatory peptides and functions in antigen processing and the cleavage of chemokines such as MIP-1. CD13 serves

as the cellular receptor for Coronavirus.

**Form:** Liquid

**Isotype:** Mouse IgG1,  $\kappa$

**Size:** 25  $\mu$ g/100  $\mu$ g

**Host:** Mouse

**Reactivity:** Human

**Application:** FCM

**Concentration:** 0.5 mg/mL

**Conjugation:** Unconjugated

**SwissProt:** P15144

**Buffer:** PBS with 0.05% Proclin300

### **Recommended Use:**

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is  $\leq 2.0 \mu$ g per  $10^6$  cells in 100  $\mu$ l volume or 100  $\mu$ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

### **Storage:**

Keep as concentrated solution.

Store at 2~8°C. Do not freeze.



This product is guaranteed up to one year from purchase.