

Manufacturer/Supplier:			
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD			
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District			
Wuxi Jiangsu Province, China			
TELE: 86-510-82732223 FAX: 86-510-82720101-8014			
Web : <u>www.dldevelop.com</u>			
Email: info@dldevelop.com.cn service@dldevelop.com.cn			
Product name:	oduct name: Anti-Human CD13 Monoclonal Antibody(PerCP/Cyanine5.5 Conjugated)		
Catalog:	DL20891F		
Synonyms:	Aminopeptidase N,ANPEP,AP-N,hAPN,Alanyl aminopeptidase,Aminopeptidase M, AP-M,Microsomal aminopeptidase,gp150,CD13		
Background:	CD13 is a 150-170 kD type II transmembrane		
	glycoprotein also known as aminopeptidase N, APN,		
	and gp150. This zinc metallopeptidase is expressed		
as a homodimer on granulocytes, myeloid			
	progenitors, endothelial cells, epithelial cells and		
	subset of granular lymphoid cells. It is not expressed		
	on platelets or erythrocytes. CD13 is thought to be		
	involved in the metabolism of many regulatory		
	peptides and functions in antigen processing and the		



cleavage of chemokines such as MIP-1. CD13 serves as the cellular receptor for Coronavirus.

Form: Liquid		Isotype: Mouse IgG1, ĸ		
Size:		Host: Mouse		
20Tests/100Te	sts/100Tests×2			
Reactivity: Human		Application: FCM		
Concentration: 5 µL		<b>Conjugation:</b> PerCP/Cyanine5.5		
SwissProt:	P15144			
Buffer:	Phosphate but	Phosphate buffered solution, pH 7.2, containing		
	0.09% stabiliz	0.09% stabilizer and 1% protein protectant.		

## **Recommended Use:**


Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

## Data:

Human peripheral blood granulocytes are stained with Anti-Human



CD13 Monoclonal Antibody(PerCP/Cyanine5.5 Conjugated)(filled gray histogram). Unstained granulocytes (empty black histogram) are used as control.



## Storage:

Keep as concentrated solution.

Store at  $2\sim 8^{\circ}$ C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.

