

## Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Human CD10 Monoclonal Antibody(Biotin

Conjugated)

**Catalog:** DL20904F

**Synonyms:** Neprilysin, MME, CALLA, NEP, Neutral

endopeptidase, SFE, CD10

**Background:** CD10 is a 100 kD neutral endopeptidase and a

> member of the metalloprotease family. It is a type II transmembrane protein also known as common acute

lymphoblastic leukemia antigen (CALLA),

enkephalinase, and neprilysin. CD10 is expressed on B cell precursors, T cell precursors, and neutrophils. CD10 is involved in B cell development and has been shown to bind opioid enkephalins, bradykinin, angiotensins I and II, and other biologically active

peptides.



Form: Liquid **Isotype:** Mouse IgG1, κ

**Size:** 25µg/100µg Host: Mouse

**Reactivity:** Human **Application:** FCM

**Concentration:** 0.5 mg/mL **Conjugation:** Biotin

SwissProt: P08473

Phosphate buffered solution, pH 7.2, containing Buffer:

0.09% stabilizer and 1% protein protectant.

## Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is  $\leq 1.0 \,\mu g$  per  $10^6$  cells in  $100 \,\mu L$  volume or  $100 \,\mu L$  of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

## **Storage:**

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do



not freeze.

This product is guaranteed up to one year from purchase.