

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road, Liangxi District
Wuxi Jiangsu Province, China

TELE: 86-510-82732223

FAX: 86-510-82720101-8014

Web : www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Human CD15 Monoclonal Antibody(PerCP
Conjugated)

Catalog: DL20917F

Synonyms: Alpha-(1,3)-fucosyltransferase 4, FUT4, ELAM-1
ligand fucosyltransferase, Fucosyltransferase
4, Fucosyltransferase IV,
Fuc-TIV, FucT-IV, Galactoside
3-L-fucosyltransferase

Background: CD15 is 3-fucosyl-N-acetylglucosamine (3-FAL),
also known as Lewis X, 3-FAL, X-hapten, and
SSEA-1. CD15 is expressed on granulocytes and
monocytes. It has also been shown to be expressed
on Langerhans cells and some malignant cells. CD15
has been implicated in adhesion, as well as
chemotaxis, phagocytosis, and bactericidal activity.

Form: Liquid

Isotype: Mouse IgM

Size:

20Tests/100Tests/100Tests×2

Host: Mouse

Reactivity: Human

Application: FCM

Concentration: 5 μ L

Conjugation: PerCP

SwissProt: P22083

Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Storage:

Keep as concentrated solution.



Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.