

Manufacturer/Supplier:						
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD						
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District						
Wuxi Jiangsu Province, China						
TELE: 86-510-82732223 FAX: 86-510-82720101-8014						
Web : <u>www.dldevelop.com</u>						
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>						
Product name:	Anti-Human CD235 Monoclonal Antibody(AF488					
	Conjugated)					
Catalog:	DL20931F					
Synonyms:	Glycophorin-A/B,GYPA/B,MN					
	sialoglycoprotein,SS-active sialoglycoprotein,PAS-2/3,Sialoglycoprotein					
alpha/delta,CD235a/b						
Background:	ackground: The HIR2 antibody reacts with a common epitope of					
glycophorin A (CD235a) and glycophorin B (CD235b). Glycophorin A is the major sialoglycoprotein expressed on red blood cell membrane, and erythroid precursors. Glycophorin A						
				shares strong homology with glycophorin B. The HIR2 antibody recognizes human RBCs and		
				erythroid precursors and is useful in erythroid cell		



development studies. Mature, non-nucleated red blood cells are characteristically glycophorin A positive, but CD45 and CD71 negative.

Form: Liquid		Isotype: Mouse IgG2b, κ	
Size:		Host: Mouse	
20Tests/100Te	sts/100Tests×2		
Reactivity: Human		Application: FCM	
Concentration: 5 µL		Conjugation: AF488	
SwissProt:	P02724,P06028		
Buffer:	Phosphate buffe	Phosphate buffered solution, pH 7.2, containing	

Recommended Use:


Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

0.09% stabilizer and 1% protein protectant.

Data:



Human peripheral blood red blood cells are stained with Anti-Human CD235 Monoclonal Antibody(AF488 Conjugated)(filled gray histogram). Unstained red blood cells (empty black histogram) are used as control.



Storage:

Keep as concentrated solution.

Store at $2\sim 8^{\circ}$ C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.