

Manufaatuvar/Si	upplice		
Manufacturer/Supplier:			
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD			
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District			
Wuxi Jiangsu Province, China			
TELE: 86-510-82732223 FAX: 86-510-82720101-801			
Web : <u>www.dldevelop.com</u>			
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>			
Product name:	Anti-Human CD33 Monoclonal Antibody(Biotin		
	Conjugated)		
Catalog:	DL20969F		
Synonyms:	Myeloid cell surface antigen CD33,CD33,Sialic		
	acid-binding Ig-like lectin 3,Siglec-3,gp67,SIGLEC3		
Background:	CD33 is a 67 kD type I transmembrane glycoprotein also known as Siglec-3, gp67, and p67. It is a		
	sialoadhesion immunoglobulin superfamily member		
	expressed on myeloid progenitors, monocytes, granulocytes, dendritic cells and mast cells. CD33 is		
	absent on normal platelets, lymphocytes,		
	erythrocytes and hematopoietic stem cells. CD33		
	functions as a sialic acid- dependent cell adhesion		
	molecule with carbohydrate/lectin binding activity.		



Form: Liquid		<b>Isotype:</b> Mouse IgG1, κ
<b>Size:</b> 25 μg/100 μg		Host: Mouse
Reactivity: Human		Application: FCM
Concentration: 0.5 mg/mL		Conjugation: Biotin
SwissProt:	P20138	
Buffer:	PBS with 0.05% Proclin300, 1% BSA	

## **Recommended Use:**

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is  $\leq 2.0 \ \mu g$  per  $10^6$  cells in 100  $\mu$ l volume or 100  $\mu$ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

## **Storage:**

Keep as concentrated solution.

Store at  $2\sim 8^{\circ}$ C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.