

**Manufacturer/Supplier:**

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

**Address:** A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District  
Wuxi Jiangsu Province, China

TELE: 86-510-82732223

FAX: 86-510-82720101-8014

**Web :** [www.dldevelop.com](http://www.dldevelop.com)

Email: [info@dldevelop.com.cn](mailto:info@dldevelop.com.cn)    [service@dldevelop.com.cn](mailto:service@dldevelop.com.cn)

**Product name:** Purified Anti-Human CD41 Monoclonal Antibody

**Catalog:** DL20985F

**Synonyms:** Integrin alpha-Iib, ITGA2B, GPalpha  
Iib, GPIIb, Platelet membrane glycoprotein  
Iib, CD41, GP2B, ITGAB

**Background:** CD41 is a 125/25 kD  $\alpha$  subunit of the gpIIb/IIIa (CD41/CD61) complex. CD41 is a heterodimer composed of a heavy chain (gpIIb $\alpha$ ) and light chain (gpIIb $\beta$ ) linked by a single disulfide bond. It is a member of the integrin family primarily expressed on platelets and megakaryocytes. CD41 has been reported to be involved with platelet aggregation and platelet attachment to the ECM. CD41/CD61 complex acts as the receptor for fibrinogen, fibronectin, Von Willebrand factor and thrombin.

**Form:** Liquid

**Isotype:** Mouse IgG1,  $\kappa$

**Size:** 25 $\mu$ g/100 $\mu$ g

**Host:** Mouse

**Reactivity:** Human

**Application:** FCM,IHC

**Concentration:** 0.5 mg/mL

**Conjugation:** Unconjugated

**SwissProt:** P08514

**Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

### **Recommended Use:**

Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is  $\leq 2.0 \mu\text{g}$  per  $10^6$  cells in 100  $\mu\text{L}$  volume or 100  $\mu\text{L}$  of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

### **Storage:**

Keep as concentrated solution.

Store at 2~8°C. Do not freeze.



This product is guaranteed up to one year from purchase.