

## Manufacturer/Supplier: WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District Wuxi Jiangsu Province, China TELE: 86-510-82732223 FAX: 86-510-82720101-8014 Web : www.dldevelop.com Email: info@dldevelop.com.cn service@dldevelop.com.cn Product name: Anti-Mouse CD103 Monoclonal Antibody(PE/Cyanine7 Conjugated) **Catalog:** DL21024F **Synonyms:** Integrin alpha-E,Itgae,Integrin alpha M290,CD103 **Background:** CD103 is a type I transmembrane glycoprotein known as aE integrin or Integrin aIEL chain. It belongs to the integrin family and is primarily found on intestinal intraepithelial lymphocytes (IEL). CD103 is also expressed on a subpopulation of lamina propria T cells, epithelial dendritic cells, lamina propria-derived dendritic cells, and a small subset of peripheral lymphocytes. T regulatory cells express high level of CD103. The CD103 expression on lymphocytes can be induced upon activation and

TGF- $\beta$  stimulation. In association with integrin  $\beta$ 7,



|                  | CD103 is express  | ed as $\alpha E/\beta 7$ heterodimer. Mature |
|------------------|---|--|
|                  | CD103 protein ca  | in be cleaved into 2 chains, a 150           |
|                  | kD (C-terminal) chain and a 25 kD (N-terminal) chain, which remain linked by disulfide bonds. |  |
|                  |   |  |
|                  | CD103 binds to E  | E-cadherin and mediates homing of            |
|                  | lymphocytes to the intestinal epithelium.   |  |
| Form: Liquid     |   | <b>Isotype:</b> Rat IgG2a, κ                 |
| Size: 25µg/100µg | į   | Host: Rat                                    |
| Reactivity: Mous | se  | Application: FCM                             |
| Concentration: ( | 0.2 mg/mL   | Conjugation: PE/Cyanine7                     |
| SwissProt:       | Q60677  |  |
| Buffer:          | Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.   |  |

## **Recommended Use:**

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to



obtain optimal results [The recommended concentration is 0.1-1  $\mu$ g/10<sup>6</sup>cells in 100  $\mu$ L volume].

## **Storage:**

Keep as concentrated solution.

Store at  $2\sim 8^{\circ}$ C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.

