

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Mouse CD117 Monoclonal Antibody(FITC

Conjugated)

Catalog: DL21053F

Synonyms: Mast/stem cell growth factor receptor

Kit, Kit, SCFR, Proto-oncogene c-Kit, Tyrosine-protein

kinase Kit, CD117

Background: CD117 is a 145 kD immunoglobulin superfamily

member also known as c-Kit and stem cell factor

receptor (SCFR). It is a transmembrane

tyrosine-kinase receptor that binds the c-Kit ligand (also known as steel factor, stem cell factor, and mast

cell growth factor). CD117 is expressed on

hematopoietic stem cells (including multipotent hematopoietic stem cells, progenitors committed to myeloid and/or erythroid lineages, and T and B cell



precursors), mast cells, and acute myeloid leukemia (AML) cells. CD117 interaction with its ligand is critical for the development of hematopoietic stem cells

Form: Liquid **Isotype:** Rat IgG2b, κ

Size: Host: Rat

50Tests/100Tests/100Tests×2

Reactivity: Mouse **Application:** FCM

Concentration: 5 µL **Conjugation: FITC**

SwissProt: P05532

Buffer: Phosphate buffered solution, pH 7.2, containing

0.09% stabilizer and 1% protein protectant.

Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.



Storage:

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.