

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Mouse CD71 Monoclonal Antibody(FITC

Conjugated)

Catalog: DL21062F

Synonyms: Transferrin receptor protein

1,Tfrc,TR,TfR,TfR1,Trfr,CD71

Background: CD71 is a 95 kD type II heterodimeric

transmembrane glycoprotein that is also known as T9

and transferrin receptor. CD71 is expressed on proliferating cells, reticulocytes, and erythroid precursors. Its expression is very low on resting leukocytes. CD71 plays a role in the control of

cellular proliferation by facilitating the uptake of iron

via ferrotransferrin binding and the recycling of

apotransferrin to the cell surface.



Form: Liquid **Isotype:** Rat IgG2a, κ

Size: Host: Rat

50Tests/100Tests/100Tests×2

Reactivity: Mouse **Application:** FCM

Concentration: 5 µL **Conjugation: FITC**

SwissProt: O62351

Phosphate buffered solution, pH 7.2, containing **Buffer:**

0.09% stabilizer and 1% protein protectant.


Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use

Data:

C57BL/6 murine bone marrow cells are stained with Anti-Mouse CD71 Monoclonal Antibody(FITC Conjugated)(filled gray histogram). Unstained bone marrow cells (empty black histogram) are used as control.





Storage:

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.