

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>

Product name: Anti-Mouse CD4 Monoclonal Antibody(PerCP

Conjugated)

Catalog: DL21135F

Synonyms: T-cell surface glycoprotein CD4,CD4,T-cell surface

antigen T4/Leu-3,CD4

Background: CD4 is a 55 kD protein also known as L3T4 or T4. It

is a member of the Ig superfamily, primarily

expressed on most thymocytes, a subset of T cells, and weakly on macrophages and dendritic cells. It acts as a coreceptor with the TCR during T cell activation and thymic differentiation by binding

MHC class II and associating with the protein tyrosin

kinase, lck.

Form: Liquid **Isotype:** Rat IgG2b, κ



Size: Host: Rat

50Tests/100Tests/100Tests×2

Reactivity: Mouse **Application:** FCM

Concentration: 5 µL **Conjugation:** PerCP

SwissProt: P06332

Buffer: Phosphate buffered solution, pH 7.2, containing

0.09% stabilizer and 1% protein protectant.


Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 µL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data:

Mouse splenocytes are stained with Anti-Mouse CD4 Monoclonal Antibody(PercP Conjugated)(filled gray histogram). Unstained splenocytes (blank black histogram) are used as control.





Storage:

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.