

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road, Liangxi District
Wuxi Jiangsu Province, China

TELE: 86-510-82732223

FAX: 86-510-82720101-8014

Web : www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Rat CD8a Monoclonal
Antibody(PerCP/Cyanine5.5 Conjugated)

Catalog: DL21162F

Synonyms: T-cell surface glycoprotein CD8 alpha
chain, CD8A, T-lymphocyte differentiation antigen
T8/Leu-2, MAL

Background: CD8a is a 32 kD glycoprotein also known as T8, Lyt2, Ly-2, and CD8 α . CD8a is a member of the immunoglobulin superfamily expressed on most thymocytes, subset of mature T cells, most NK cells, macrophages, and some activated CD4⁺ T cells (not resting). CD8a forms heterodimers with the CD8 β chain (CD8b) on the surface of most thymocytes, while mature peripheral T lymphocytes express almost exclusively the CD8 $\alpha\beta$ heterodimer.

Intestinal intraepithelial lymphocytes express CD8a without CD8b. CD8 is an antigen co-receptor on T cells that interacts with MHC class I on antigen-presenting cells or epithelial cells. CD8 participates in T cell activation through its association with the T cell receptor complex and protein tyrosine kinase lck (p56lck).

Form: Liquid

Isotype: Mouse IgG1, κ

Size:

Host: Mouse

50Tests/100Tests/200Tests/500Tests

Reactivity: Rat

Application: FCM

Concentration:

Conjugation: PerCP/Cyanine5.5

SwissProt: P07725

Buffer: PBS with 0.05% Proclin300, 1% BSA

Recommended Use:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million

cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data:

Rat splenocytes are stained with Anti-Rat CD8a Monoclonal Antibody(PerCP/Cyanine5.5 Conjugated)(filled gray histogram)or Mouse IgG1, κ Isotype Control(PerCP/Cyanine5.5 Conjugated)(empty black histogram).

Storage:

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.