

Manufacturer/Supplier:	
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD	
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District	
Wuxi Jiangsu Province, China	
TELE: 86-510-8	2732223 FAX: 86-510-82720101-8014
Web : <u>www.dldevelop.com</u>	
Email: info@dldevelop.com.cn service@dldevelop.com.cn	
Product name:	Anti-Mouse IFN-γ Monoclonal Antibody(APC
	Conjugated)
Catalog:	DL21201F
c	222.2011
Synonyms:	Interferon gamma,Ifng,IFN-gamma, IFNγ
Synonyms.	interferon gamma, ing, in iv gamma, in ivy
Background:	IFN- γ is a potent multifunctional cytokine which is
Dackground.	secreted primarily by activated NK cells and T cells.
	Originally characterized based on anti-viral activities,
	IFN-γ also exerts anti-proliferative,
	immunoregulatory, and proinflammatory activities.
	IFN-γ can upregulate MHC class I and II antigen
	expression by antigen-presenting cells.
- - · · · · · · · · · · · · · · · · · · ·	
Form: Liquid	Isotype: Rat IgG1, ĸ
<u>C:</u>	Harte D. (
Size:	Host: Rat



50Tests/100Tests/100Tests×2 Reactivity: Mouse

Application: ICFCM

Concentration: 5 µL

Conjugation: APC

SwissProt: P01580

Buffer:Phosphate buffered solution, pH 7.2, containing
0.09% stabilizer and 1% protein protectant.

Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Storage:

Keep as concentrated solution.

Store at $2\sim 8^{\circ}$ C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.