

## Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

**Web**: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Mouse CD25 Monoclonal

Antibody(PerCP/Cyanine5.5 Conjugated)

**Catalog:** DL21223F

**Synonyms:** Interleukin-2 receptor subunit alpha, IL2RA, IL-2

receptor subunit alpha, IL-2-RA, IL-2R subunit

alpha,IL2-RA,TAC antigen,p55,p55

**Background:** CD25 is a 55 kD glycoprotein, also known as the low

> affinity IL-2Rα, Ly-43, p55, or Tac. It is expressed on activated T and B cells, thymocyte subset, pre-B cells, and T regulatory cells. In association with CD122 (IL-2Rβ) and CD132(common γ chain), CD25 forms the high affinity signaling IL-2 receptor.

Form: Liquid **Isotype:** Rat IgG1, κ



**Size:** 25µg/100µg Host: Rat

**Reactivity:** Mouse **Application:** FCM

**Concentration:** 0.2 mg/mL Conjugation: PerCP/Cyanine5.5

**SwissProt:** P01590

Buffer: Phosphate buffered solution, pH 7.2, containing

0.09% stabilizer and 1% protein protectant.


## Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1  $\mu g/10^6$  cells in 100  $\mu L$  volume].

## Data:

C57BL/6 murine splenocytes are stained with Anti-Mouse CD25 Monoclonal Antibody(PerCP/Cyanine 5.5 Conjugated)[Used at 0.2 ug/10<sup>6</sup> cells dilution] and Anti-Mouse CD4 Monoclonal Antibody (FITC Conjugated).





## **Storage:**

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.