

## Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: 

Antibody(PE/Cyanine5.5 Conjugated)

**Catalog:** DL21234F

**Synonyms:** T-cell surface glycoprotein CD3 epsilon

chain, CD3E, T-cell surface antigen T3/Leu-4 epsilon

chain,CD3e,CD3E,T3E

**Background:** CD3\varepsilon is a 20 kD transmembrane protein, also known

> as CD3 or T3. It is a member of the Ig superfamily and primarily expressed on T cells, NK-T cells, and at different levels on thymocytes during T cell differentiation. CD3E forms a TCR complex by associating with the CD3 $\delta$ ,  $\gamma$  and  $\zeta$  chains, as well as the TCR  $\alpha/\beta$  or  $\gamma/\delta$  chains. CD3 plays a critical role in TCR signal transduction, T cell activation, and

antigen recognition by binding the peptide/MHC



## antigen complex.

Form: Liquid Isotype: Armenian Hamster IgG

Size: Host: Armenian Hamster

50Tests/100Tests/100Tests×2

Reactivity: Mouse Application: FCM

Concentration: 5 µL Conjugation: PE/Cyanine5.5

SwissProt: P22646

**Buffer:** Phosphate buffered solution, pH 7.2, containing

0.09% stabilizer and 1% protein protectant.

## **Recommended Use:**

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

## Storage:



Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.