

| Manufacturer/Si | unlier | | |
|--------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------|--|--|
| WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD | | | |
| Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District | | | |
| Wuxi Jiangsu Province, China | | | |
| | | | |
| TELE: 86-510-82732223 FAX: 86-510-82720101-8014 | | | |
| Web : <u>www.dldevelop.com</u> | | | |
| Email: info@dldevelop.com.cn service@dldevelop.com.cn | | | |
| | | | |
| Product name: | Anti-Mouse CD8a Monoclonal | | |
| | Antibody(PE/Cyanine5.5 Conjugated) | | |
| Catalog: | DL21266F | | |
| _ | | | |
| Synonyms: | T-cell surface glycoprotein CD8 alpha chain,CD8A,T-lymphocyte differentiation antigen | | |
| | | | |
| | T8/Leu-2,MAL | | |
| | | | |
| Background: | Background: CD8, also known as Lyt-2, Ly-2, or T8, consists of disulfide-linked α and β chains that form the | | |
| | | | |
| | α (CD8a)/ β (CD8b) heterodimer and α/α homodimer. CD8a is a 34 kD protein that belongs to the | | |
| | | | |
| | immunoglobulin family. The CD8 α/β heterodimer is | | |
| | expressed on the surface of most thymocytes and a | | |
| | subset of mature TCR α/β T cells. CD8 expression on | | |
| | mature T cells is non- overlapping with CD4. The | | |
| | | | |
| | CD8 α/α homodimer is expressed on a subset of γ/δ | | |
| | | | |



| | TCR-bearing T cells, NK cells, intestinal intraepithelial lymphocytes, and lymphoid dendritic cells. CD8 is an antigen co-receptor on T cells that interacts with MHC class I on antigen-presenting cells or epithelial cells. CD8 promotes T cell activation through its association with the TCR complex and protein tyrosine kinase lck. | | |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------|--|
| Form: Liquid | | Isotype: Rat IgG2a, κ | |
| Size: 25µg/100µg | | Host: Rat | |
| Reactivity: Mouse | | Application: FCM | |
| Concentration: 0.2 mg/mL | | Conjugation: PE/Cyanine5.5 | |
| SwissProt: | P01731 | | |
| Buffer: | Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant. | | |

Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for



individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 μ g/10⁶cells in 100 μ L volume].

Storage:

Keep as concentrated solution.

Store at $2\sim 8^{\circ}$ C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.

