

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Rat CD4(domain 1) Monoclonal

Antibody(PerCP Conjugated)

Catalog: DL21285F

Synonyms: T-cell surface glycoprotein CD4,CD4,T-cell surface

antigen T4/Leu-3,CD4

Background: CD4, also known as T4, is a 55kD glycoprotein

member of the immunoglobin superfamily and is expressed on majority of thymocytes, macrophages, and a peripheral T cell subset (T helper cells). CD4 is a T cell co-receptor that interacts with the MHC class

II molecule and is involved in T cell activation.

Form: Liquid **Isotype:** Mouse IgG2a, κ

Size: 25μg/100μg Host: Mouse



Reactivity: Rat **Application:** FCM

Concentration: 0.2 mg/mL **Conjugation:** PerCP

SwissProt: P05540

Phosphate buffered solution, pH 7.2, containing Buffer:

0.09% stabilizer and 1% protein protectant.


Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 $\mu g/10^6$ cells in 100 μL volume].

Data:

Rat splenocytes are stained with Anti-Rat CD4(domain 1 Conjugated) Monoclonal Antibody (PerCP Conjugated) [Used at 0.1 μg/10⁶ cells dilution](filled gray histogram). Unstained splenocytes (empty black histogram) are used as control.





Storage:

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.