

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Purified Anti-Human CD4 Monoclonal Antibody

Catalog: DL21322F

Synonyms: T-cell surface glycoprotein CD4,CD4,T-cell surface

antigen T4/Leu-3,CD4

Background: CD4, also known as T4/Leu-3, is a 55 kD

> single-chain type I transmembrane glycoprotein and member of the immunoglobulin superfamily. It is expressed on most thymocytes, helper T cells, type II NKT cells, and monocytes/macrophages. CD4 is part of the TCR/CD3 complex, binds to β2 domain from the MHC class II molecule, and participates in TCR signal transduction. CD4 is the receptor of IL-16 and is a coreceptor for the human immunodeficiency virus (HIV) and human herpes virus 7 (HHV-7).



Form: Liquid **Isotype:** Mouse IgG1, κ

Size: 25µg/100µg Host: Mouse

Reactivity: Human **Application:** FCM,IHC,IF

Concentration: 0.5 mg/mL Conjugation: Unconjugated

SwissProt: P01730

Buffer: Phosphate buffered solution, pH 7.2, containing

0.09% stabilizer

Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is $\leq 2.0 \mu g$ per 10^6 cells in $100 \mu L$ volume or $100 \mu L$ of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Storage:

Keep as concentrated solution.

Store at $2\sim8^{\circ}$ C. Do not freeze.

This product is guaranteed up to one year from purchase.