

**Manufacturer/Supplier:**

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

**Address:** A1-203 Mingpin CityII, No.8 Xihudong Road, Liangxi District  
Wuxi Jiangsu Province, China

TELE: 86-510-82732223

FAX: 86-510-82720101-8014

**Web :** [www.dldevelop.com](http://www.dldevelop.com)

Email: [info@dldevelop.com.cn](mailto:info@dldevelop.com.cn)    [service@dldevelop.com.cn](mailto:service@dldevelop.com.cn)

**Product name:** Anti-Human CD4 Monoclonal Antibody(AF488  
Conjugated)

**Catalog:** DL21332F

**Synonyms:** T-cell surface glycoprotein CD4, CD4, T-cell surface  
antigen T4/Leu-3, CD4

**Background:** CD4, also known as T4/Leu-3, is a 55 kD  
single-chain type I transmembrane glycoprotein and  
member of the immunoglobulin superfamily. It is  
expressed on most thymocytes, helper T cells, type II  
NKT cells, and monocytes/macrophages. CD4 is part  
of the TCR/CD3 complex, binds to  $\beta 2$  domain from  
the MHC class II molecule, and participates in TCR  
signal transduction. CD4 is the receptor of IL-16 and  
is a coreceptor for the human immunodeficiency  
virus (HIV) and human herpes virus 7 (HHV-7).

**Form:** Liquid

**Isotype:** Mouse IgG1,  $\kappa$

**Size:**

20Tests/100Tests/100Tests $\times$ 2

**Host:** Mouse

**Reactivity:** Human

**Application:** FCM

**Concentration:** 5  $\mu$ L

**Conjugation:** AF488

**SwissProt:** P01730

**Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.


### **Recommended Use:**

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

### **Data:**

Human peripheral blood cells are stained with Anti-Human CD4 Monoclonal Antibody(AF488 Conjugated)(filled gray histogram). Unstained peripheral blood cells (blank black histogram) are used as

control.



### Storage:

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.