

## Manufacturer/Supplier: WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District Wuxi Jiangsu Province, China TELE: 86-510-82732223 FAX: 86-510-82720101-8014 Web : www.dldevelop.com Email: info@dldevelop.com.cn service@dldevelop.com.cn Product name: Anti-Mouse CD45R/B220 Monoclonal Antibody(PerCP Conjugated) **Catalog:** DL21376F **Synonyms:** B220,CD45,GP180,LCA,T200 **Background:** CD45R, also known as B220, is an isoform of CD45. It is a member of the protein tyrosine phosphatase (PTP) family with a molecular weight of approximately 180-240 kD. CD45R is expressed on B cells (at all developmental stages from pro-B cells through mature B cells), activated B cells, and subsets of T and NK cells. CD45R (B220) is also expressed on a subset of abnormal T cells involved in the pathogenesis of systemic autoimmunity in MRL-Faslpr and MRL- Fasgld mice. It plays a critical role in TCR and BCR signaling. The primary



ligands for CD45 are galectin-1, CD2, CD3, and CD4. CD45R is commonly used as a pan-B cell marker; however, CD19 may be more appropriate for B cell specificity.

Form: Liquid		Isotype: Rat IgM
Size: 25µg/100µg		Host: Rat
Reactivity: Mouse		Application: FCM
Concentration: 0.2 mg/mL		Conjugation: PerCP
SwissProt:	P06800,P08575	
Buffer:	Phosphate buffered solution, pH 7.2, containing	

#### **Recommended Use:**


Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1  $\mu g/10^6$ cells in 100  $\mu$ L volume].

0.09% stabilizer and 1% protein protectant.

# **DEVELOP**.

### Data:

C57BL/6 murine splenocytes are stained with Anti-Mouse CD45R/B220 Monoclonal Antibody(PerCP Conjugated)[Used at 0.05  $\mu$ g/10<sup>6</sup> cells dilution](filled gray histogram). Unstained splenocytes (empty black histogram) are used as control.



### **Storage:**

Keep as concentrated solution.

Store at  $2\sim 8^{\circ}$ C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.