

Manufacturer/Su	upplier:		
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD			
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District			
Wuxi Jiangsu Province, China			
TELE: 86-510-82732223 FAX: 86-510-82720101-801			
Web : <u>www.dldevelop.com</u>			
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>			
Product name:	Anti-Mouse CD183/CXCR3 Monoclonal		
	Antibody(AF488 Conjugated)		
Catalog:	DL21409F		
Synonyms:	C-X-C chemokine receptor type		
	,Cxcr3,CXC-R3,CXCR-3,Interferon-inducible		
	protein 10 receptor, IP-10 receptor, CD183/CXCR3		
Background:	CD183/CXCR3, also known as CXCR3, is a member		
of the C-X-C chemokine family, characterized by			
pair of cysteine residues separated by a single amino			
acid. CXCR3 is a 38 kD seven pass transmembrane			
receptor coupled to G-protein. It mediates Ca2+			
mobilization and chemotaxis in response to C-X-C			
chemokines, such as IP10 (CXCL10), MIG			
(CXCL9), I-TAC (CXCL11) and PF4 (CXCL4). CXCR3 is expressed primarily on activiated T			



lymphocytes, NK cells, and some epithelial cells and endothelial cells. It is not expressed on B cells, monocytes or granulocytes.

Form: Liquid		Isotype: Armenian Hamster IgG
Size: 25µg/100µg		Host: Armenian Hamster
Reactivity: Mouse		Application: FCM
Concentration: 0.5 mg/mL		Conjugation: AF488
SwissProt:	O88410	
Buffer:	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.	


Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 $\mu g/10^6$ cells in 100 μ L volume].

Data:



C57BL/6 murine splenocytes are stained with Anti-Mouse CD183/CXCR3 Monoclonal Antibody (AF488 Conjugated)[Used at 0.2 μ g/10⁶ cells dilution](filled gray histogram). Unstained splenocytes (empty black histogram) are used as control.



Storage:

Keep as concentrated solution.

Store at $2\sim 8^{\circ}$ C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.