

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Mouse CD49b Monoclonal Antibody(FITC

Conjugated)

Catalog: DL21420F

Synonyms: Integrin alpha-2,CD49 antigen-like family member

B, Collagen receptor, Platelet membrane glycoprotein

Ia, GPIa, VLA-2 subunit alpha, CD49b

Background: DX5 antigen has been recently characterized as

> CD49b. It is a 150 kD integrin α chain also known as α 2 integrin, VLA-2 α chain, and integrin α 2 chain. CD49b non-covalently associates with CD29 (\beta1 integrin) to form the CD49b/CD29 complex known as VLA-2, a receptor for collagen and laminin.

CD49b is expressed on platelets, the majority of NK cells, NKT cells, and a small subset of CD8+ T cells

(this population can be significantly increased



following viral infection). DX5 is used for the identification and isolation of NK cells, and is especially useful for identifying NK cells in mice lacking the NK1.1 antigen.

Form: Liquid **Isotype:** Rat IgM

Size: Host: Rat

50Tests/100Tests/100Tests×2

Reactivity: Mouse **Application:** FCM

Concentration: 5 µL **Conjugation:** FITC

SwissProt: O62469

Buffer: Phosphate buffered solution, pH 7.2, containing

0.09% stabilizer and 1% protein protectant.


Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.



Data:

Mouse bone marrow cells are stained with Anti-Mouse CD49b Monoclonal Antibody(FITC Conjugated)(filled gray histogram). Unstained bone marrow cells (blank black histogram) are used as control.



Storage:

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.