

## Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

**Web**: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Human CD123 Monoclonal Antibody(FITC

Conjugated)

**Catalog:** DL21442F

**Synonyms:** IL-3Rα, IL-3 Receptor alpha

**Background:** CD123 is the 70 kD transmembrane  $\alpha$  chain of the

> IL-3 receptor. Alone, CD123 binds IL-3 with low affinity; when CD123 associates with CDw131 (common  $\beta$  chain), it binds IL-3 with high affinity. CD123 does not transduce intracellular signals upon

binding IL-3 and requires the  $\beta$  chain for this

function. CD123 is expressed by myeloid precursors, macrophages, dendritic cells, mast cells, basophils,

megakaryocytes, and some B cells.

Form: Liquid **Isotype:** Mouse IgG1, κ



Size: 20 Tests/50 Tests/100 Host: Mouse

Tests/200 Tests

**Reactivity:** Human **Application:** FCM

**Concentration: Conjugation: FITC** 

SwissProt: P26951

Buffer: PBS with 0.05% Proclin300, 1% BSA

## Recommended Use:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 µL staining volume or per 100 µL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

## **Storage:**

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.