

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road, Liangxi District
Wuxi Jiangsu Province, China

TELE: 86-510-82732223

FAX: 86-510-82720101-8014

Web : www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Human CD14 Monoclonal Antibody(Biotin
Conjugated)

Catalog: DL21462F

Synonyms: Monocyte differentiation antigen
CD14, CD14, Myeloid cell-specific leucine-rich
glycoprotein

Background: CD14 is a 53-55 kD glycosylphosphatidylinositol (GPI)-linked membrane glycoprotein that is also known as the LPS receptor. CD14 is expressed at high levels on monocytes and macrophages, and at lower levels on granulocytes. Some dendritic cell populations such as interfollicular dendritic cells, reticular dendritic cells, and Langerhans cells have also been reported to express CD14. As a high-affinity receptor for LPS, CD14 is involved in

the clearance of gram-negative pathogens and in the upregulation of adhesion molecules and cytokine expression in monocytes and neutrophils.

Form: Liquid

Isotype: Mouse IgG1, κ

Size: 25 μ g/100 μ g

Host: Mouse

Reactivity: Human

Application: FCM

Concentration: 0.5 mg/mL

Conjugation: Biotin

SwissProt: P08571

Buffer: PBS with 0.05% Proclin300, 1% BSA

Recommended Use:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is $\leq 0.5 \mu$ g per 10^6 cells in 100 μ l volume or 100 μ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Storage:



Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.