

Manufacturer/Supplier:			
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD			
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District			
Wuxi Jiangsu Pro	vince, China		
TELE: 86-510-82732223 FAX: 86-510-82720101-80			
Web : www.dldev	velop.com		
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>			
Product name:	Anti-Mouse CD45 Monoclonal		
	Antibody(PE/Cyanine5 Conjugated)		
Catalog:	DL21681F		
Synonyms:	Receptor-type tyrosine-protein phosphatase		
	C,Ptprc,L-CA,Ly-5,T200,CD45		
Background:	ckground: CD45 is a 180-240 kD glycoprotein also known as		
the leukocyte common antigen (LCA), T200, or Ly-5. It is a member of the protein tyrosine			
			phosphatase (PTP) family, expressed on all
	hematopoietic cells except mature erythrocytes and		
	platelets. There are different isoforms of CD45 that		
	arise from alternative splicing of exons 4, 5, and 6,		
	which encode A, B, and C determinants,		
	respectively. CD45 plays a key role in TCR and BCR		
	signal transduction. These isoforms are very specific		



to the activation and maturation state of the cell as well as cell type. The primary ligands for CD45 are galectin-1, CD2, CD3, CD4, TCR, CD22, and Thy-1.

Form: Liquid		Isotype: Rat IgG2b, κ	
Size:		Host: Rat	
50Tests/100Te	sts/100Tests×2		
Reactivity: Mouse		Application: FCM	
Concentration: 5 µL		Conjugation: PE/Cyanine5	
SwissProt:	P06800		
Buffer:	Phosphate bu	Phosphate buffered solution, pH 7.2, containing	

Recommended Use:


Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

0.09% stabilizer and 1% protein protectant.

Data:



C57BL/6 murine splenocytes are stained with Anti-Mouse CD45 Monoclonal Antibody(PE/Cyanine5 Conjugated)(filled gray histogram). Unstained splenocytes (empty black histogram) are used as control.



Storage:

Keep as concentrated solution.

Store at $2\sim 8^{\circ}$ C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.

