

Manufacturer/Supplier:			
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD			
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District			
Wuxi Jiangsu Province, China			
TELE: 86-510-82732223   FAX: 86-510-82720101-8014			
Web : <u>www.dldevelop.com</u>			
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>			
Product name:	Anti-Mouse CD45 Monoclonal Antibody(AF488		
	Conjugated)		
Catalog:	DL21695F		
Synonyms:	Receptor-type tyrosine-protein phosphatase		
	C,Ptprc,L-CA,Ly-5,T200,CD45		
<b>Background:</b>	CD45 is a 180-240 kD glycoprotein also known as		
	the leukocyte common antigen (LCA), T200, or		
Ly-5. It is a member of the protein tyrosine			
phosphatase (PTP) family, expressed on all			
hematopoietic cells except mature erythrocytes and			
platelets. There are different isoforms of CD45 that			
	arise from alternative splicing of exons 4, 5, and 6,		
	which encode A, B, and C determinants,		
	respectively. CD45 plays a key role in TCR and BCR		
	signal transduction. These isoforms are very specific		



to the activation and maturation state of the cell as well as cell type. The primary ligands for CD45 are galectin-1, CD2, CD3, CD4, TCR, CD22, and Thy-1.

Form: Liquid		Isotype: Rat IgG2b, ĸ
Size: 25µg/100µg		Host: Rat
Reactivity: Mouse		Application: FCM
Concentration: 0.5 mg/mL		Conjugation: AF488
SwissProt:	P06800	

Buffer:Phosphate buffered solution, pH 7.2, containing<br/>0.09% stabilizer and 1% protein protectant.


## **Recommended Use:**

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1  $\mu$ g/10<sup>6</sup>cells in 100  $\mu$ L volume].

## Data:



C57BL/6 murine splenocytes are stained with Anti-Mouse CD45 Monoclonal Antibody(AF488 Conjugated)[Used at 0.2  $\mu$ g/10<sup>6</sup> cells dilution](filled gray histogram). Unstained splenocytes (empty black histogram) are used as control.



## **Storage:**

Keep as concentrated solution.

Store at  $2\sim 8^{\circ}$ C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.