

Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82732223 FAX: 86-510-82720101-8014				
Web : <u>www.dldevelop.com</u>				
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>				
Product name:	Anti-Human CD45 Monoclonal Antibody(PE			
	Conjugated)			
Catalog:	DL21698F			
Synonyms:	Receptor-type tyrosine-protein phosphatase			
	C,Ptprc,L-CA,Ly-5,T200,CD45			
<b>Background:</b>	CD45 is a 180 - 240 kD single chain type I			
membrane glycoprotein also known as leukocyte				
common antigen (LCA) and T200. It is a tyrosine				
phosphatase expressed on the plasma membrane of				
all hematopoietic cells, except erythrocytes or				
platelets. CD45 is a signaling molecule that regulates				
a variety of cellular processes including cell growth,				
	differentiation, cell cycle, and oncogenic			
transformation. CD45 plays a critical role in T and B				
	cell antigen receptor- mediated activation by			



dephosphorylating substrates including p56Lck,		
p59Fyn, and other Src family kinases. CD45		
non-covalently associates with lymphocyte		
phosphatase-associated phosphoprotein (LPAP) on T		
and B lymphocytes. CD45 has been reported to bind		
galectin-1 and to be associated with several other cell		
surface antigens including CD1, CD2, CD3, and		
CD4.		

Form: Liquid		<b>Isotype:</b> Mouse IgG1, κ
Size: 20Tests/100Tests/	100Tests×2	Host: Mouse
Reactivity: Human		Application: FCM
Concentration: 5 µL		Conjugation: PE
SwissProt:	P08575	
Buffer:	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.	

## **Recommended Use:**


Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5



 $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

## Data:

Human peripheral blood lymphocytes are stained with Anti-Human CD45 Monoclonal Antibody(PE Conjugated)(filled gray histogram). Unstained lymphocytes (empty black histogram) are used as control.



## **Storage:**

Keep as concentrated solution.

Store at  $2\sim 8^{\circ}$ C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.

www.dldevelop.com