

## Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Anti-Human TCR γ/δ Monoclonal Antibody(PE Product name:

Conjugated)

**Catalog:** DL21769F

T cell receptor  $\gamma/\delta$ ,  $\gamma/\delta$  TCR, TCR- $\gamma/\delta$ **Synonyms:** 

**Background:** CD49d is a 150 kD α integrin chain known as α4

> integrin or VLA-4 α chain. It forms a heterodimer with either integrin  $\beta$ 1 ( $\alpha$ 4 $\beta$ 1, VLA-4) or  $\beta$ 7 ( $\alpha$ 4 $\beta$ 7). CD49d is expressed broadly on T lymphocytes, B lymphocytes, monocytes, thymocytes, eosinophils, basophils, mast cells, NK cells, dendritic cells, and some non-hematopoietic cells, but not on normal red blood cells, platelets or neutrophils. VLA-4 binds to VCAM-1 (CD106) and fibronectin.  $\alpha 4\beta 7$  is the receptor for VCAM-1 and MAdCAM-1. CD49d

participates in mononuclear cell trafficking to



endothelial sites of inflammation and has roles in cell-cell interactions and cell adhesion to extracellular matrices. CD49d is involved in lymphocyte migration, T cell activation, and hematopoietic stem cell differentiation. CD49d is a marker to isolate pure populations of Treg cells due to its absence on Foxp3+ cells.

Form: Liquid **Isotype:** Mouse IgG1, κ

Size: 20 Tests/50 Tests/100 **Host:** Mouse

Tests/200 Tests

**Application:** FCM **Reactivity:** Human

**Concentration: Conjugation: PE** 

SwissProt:

**Buffer:** PBS with 0.05% Proclin300, 1% BSA

## Recommended Use:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 µL staining volume or per 100 µL of whole blood). Please



check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

## **Storage:**

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.