

**Manufacturer/Supplier:**

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

**Address:** A1-203 Mingpin CityII, No.8 Xihudong Road, Liangxi District  
Wuxi Jiangsu Province, China

TELE: 86-510-82732223

FAX: 86-510-82720101-8014

**Web :** [www.dldevelop.com](http://www.dldevelop.com)

Email: [info@dldevelop.com.cn](mailto:info@dldevelop.com.cn)    [service@dldevelop.com.cn](mailto:service@dldevelop.com.cn)

**Product name:** Anti-Human TCR  $\gamma/\delta$  Monoclonal Antibody(AF488  
Conjugated)

**Catalog:** DL21775F

**Synonyms:** T cell receptor  $\gamma/\delta$ ,  $\gamma/\delta$  TCR, TCR- $\gamma/\delta$

**Background:** CD49d is a 150 kD  $\alpha$  integrin chain known as  $\alpha 4$  integrin or VLA-4  $\alpha$  chain. It forms a heterodimer with either integrin  $\beta 1$  ( $\alpha 4\beta 1$ , VLA-4) or  $\beta 7$  ( $\alpha 4\beta 7$ ). CD49d is expressed broadly on T lymphocytes, B lymphocytes, monocytes, thymocytes, eosinophils, basophils, mast cells, NK cells, dendritic cells, and some non-hematopoietic cells, but not on normal red blood cells, platelets or neutrophils. VLA-4 binds to VCAM-1 (CD106) and fibronectin.  $\alpha 4\beta 7$  is the receptor for VCAM-1 and MAdCAM-1. CD49d participates in mononuclear cell trafficking to

endothelial sites of inflammation and has roles in cell-cell interactions and cell adhesion to extracellular matrices. CD49d is involved in lymphocyte migration, T cell activation, and hematopoietic stem cell differentiation. CD49d is a marker to isolate pure populations of Treg cells due to its absence on Foxp3+ cells.

**Form:** Liquid

**Isotype:** Mouse IgG1,  $\kappa$

**Size:** 20 Tests/50 Tests/100 Tests/200 Tests

**Host:** Mouse

**Reactivity:** Human

**Application:** FCM

**Concentration:**

**Conjugation:** AF488

**SwissProt:**

**Buffer:** PBS with 0.05% Proclin300, 1% BSA

### **Recommended Use:**

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please

check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

**Storage:**

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.