

M f 4/6-	1 <b>:</b>		
Manufacturer/Supplier:			
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD			
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District			
Wuxi Jiangsu Province, China			
TELE: 86-510-82732223 FAX: 86-510-82720101-8014			
Web : <u>www.dldevelop.com</u>			
Email: info@dldevelop.com.cn service@dldevelop.com.cn			
Product name:	Anti-Human TCR γ/δ Monoclonal Antibody(AF647		
	Conjugated)		
Catalog:	DL21776F		
0			
Synonyms:	T cell receptor $\gamma/\delta$ , $\gamma/\delta$ TCR, TCR- $\gamma/\delta$		
Background:	CD49d is a 150 kD α integrin chain known as α4		
	integrin or VLA-4 $\alpha$ chain. It forms a heterodimer		
	with either integrin $\beta$ 1 ( $\alpha$ 4 $\beta$ 1, VLA-4) or $\beta$ 7 ( $\alpha$ 4 $\beta$ 7).		
	CD49d is expressed broadly on T lymphocytes, B		
	lymphocytes, monocytes, thymocytes, eosinophils,		
	basophils, mast cells, NK cells, dendritic cells, and		
	some non-hematopoietic cells, but not on normal red		
	• · · · ·		
	blood cells, platelets or neutrophils. VLA-4 binds to		
	VCAM-1 (CD106) and fibronectin. $\alpha 4\beta 7$ is the		
	receptor for VCAM-1 and MAdCAM-1. CD49d		
	participates in mononuclear cell trafficking to		



κ

endothelial sites of inflammation and has roles in cell-cell interactions and cell adhesion to extracellular matrices. CD49d is involved in lymphocyte migration, T cell activation, and hematopoietic stem cell differentiation. CD49d is a marker to isolate pure populations of Treg cells due to its absence on Foxp3+ cells.

Form: Liquid	Isotype: Mouse IgG1,
Size: 20 Tests/50 Tests/100	Host: Mouse
Tests/200 Tests	
Reactivity: Human	<b>Application:</b> FCM
Concentration:	Conjugation: AF647

**SwissProt:** 

**Buffer:** PBS with 0.05% Proclin300, 1% BSA

## **Recommended Use:**

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please



check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

## **Storage:**

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.

