

## Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Human CD11b Monoclonal Antibody(PE

Conjugated)

**Catalog:** DL21778F

**Synonyms:** Integrin alpha-M,Itgam,CD11 antigen-like family

member B,CR-3 alpha chain,Leukocyte adhesion

receptor MO1,CD11b

**Background:** CD11b is a 165-170 kD type I transmembrane

> glycoprotein also known as aMintegrin, Mac-1, CR3, and C3biR. CD11b non-covalently associates with integrin β2(CD18) and is expressed on granulocytes, monocytes/macrophages, dendritic cells, NK cells, and subsets of T and B cells. CD11b/CD18 is critical for the transendothelial migration of monocytes and

neutrophils. It is also involved in granulocyte adhesion, phagocytosis, and neutrophil activation.



CD11b/CD18 interacts with ICAM-1 (CD54), ICAM-2 (CD102), ICAM-4, CD14, CD23, heparin, iC3b, fibrinogen, and factor X.

Form: Liquid **Isotype:** Mouse IgG1, κ

Size: Host: Mouse

20Tests/100Tests/100Tests×2

**Reactivity:** Human **Application:** FCM

Concentration: 5 µL Conjugation: PE

**SwissProt**: P11215

**Buffer:** Phosphate buffered solution, pH 7.2, containing

0.09% stabilizer and 1% protein protectant.

## Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.



## **Storage:**

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.