

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Human CD117 Monoclonal Antibody(FITC

Conjugated)

Catalog: DL21809F

Synonyms: Mast/stem cell growth factor receptor

Kit, Kit, SCFR, Proto-oncogene c-Kit, Tyrosine-protein

kinase Kit, CD117

Background: CD117 is a 145 kD protein tyrosine kinase also

known as c-Kit. It is a receptor for stem cell factor or

c-Kit ligand. CD117 is expressed on pluripotent hematopoietic progenitor cells (approximately 1-4% bone marrow cells), mast cells, and acute myeloid leukemia cells (AML). CD117 binding of c-Kit ligand induces phosphorylation of CD117 and stimulates proliferation and survival of primitive

hematopoietic stem cells as well as



erythroid-committed and granulo- monocytic committed cells.

Form: Liquid **Isotype:** Mouse IgG1, κ

Size: **Host:** Mouse

20Tests/100Tests/100Tests×2

Application: FCM **Reactivity:** Human

Concentration: 5 µL **Conjugation:** FITC

SwissProt: P10721

Buffer: Phosphate buffered solution, pH 7.2, containing

0.09% stabilizer and 1% protein protectant.

Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.



Storage:

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.