

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Human CD184/CXCR4 Monoclonal

Antibody(PE Conjugated)

Catalog: DL21842F

Synonyms: FB22, Fusin, HM89, LCR1, LAP-3

Background: CD184, also known as fusin or CXCR4, is a 45 kD

seven transmembrane G-protein-linked CXC

chemokine receptor. CD184 is widely expressed on

blood and tissue cells, including B and T cells,

monocytes, macrophages, dendritic cells,

granulocytes, megakaryocytes/platelets, lymphoid, myeloid precursor cells, endothelial cells, epithelial cells, astrocytes, and neurons, among other tissue cells. CD184 is the receptor for CXC chemokine SDF-1, mediates blood cell migration, and is

involved in B lymphopoiesis and myelopoiesis,



cardiogenesis, blood vessel formation, and cerebellar development. CXCR4 is also a coreceptor of X4 HIV-1 and an alternative receptor for some isolates of HIV-2.

Form: Liquid **Isotype:** Mouse IgG2a, κ

Size: Host: Mouse

20Tests/100Tests/100Tests×2

Reactivity: Human **Application:** FCM

Concentration: 5 µL Conjugation: PE

SwissProt: P61073

Buffer: Phosphate buffered solution, pH 7.2, containing

0.09% stabilizer and 1% protein protectant.

Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.



Storage:

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.