

Manufacturer/Supplier:			
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD			
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District			
Wuxi Jiangsu Province, China			
TELE: 86-510-82732223 FAX: 86-510-82720101-8014			
Web : <u>www.dldevelop.com</u>			
Email: info@dldevelop.com.cn service@dldevelop.com.cn			
Product name:	Anti-Human Ig light chain κ Monoclonal		
	Antibody(FITC Conjugated)		
Catalog:	DL21898F		
Synonyms:	Immunoglobulin light chain kappa,IGKC		
Background:	The MHK-49 antibody reacts with both soluble and		
	membrane human immunoglobulin light chain kappa		
	( $\kappa$ ). It does not react with human immunoglobulin		
	light chain lambda ( $\lambda$ ) or heavy chain. The MHK-49		
	antibody can be used as primary or secondary reagent		
	for immunofluorescent staining or ELISA analysis.		
Form: Liquid	Isotype: Mo	use IgG1, κ	
Size: 20 Tests/50	Tests/100 Host: Mouse	9	
Tests/200 Tests			



## Reactivity: Human

**Application:** FCM

**Concentration:** 

**Conjugation:** FITC

SwissProt:	P01834
SwissProt:	P01834

**Buffer:** PBS with 0.05% Proclin300, 1% BSA

## **Recommended Use:**

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

## **Storage:**

Keep as concentrated solution.

Store at  $2\sim 8^{\circ}$ C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.