

| Manufacturer/Supplier: | | |
|--|---|---|
| WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD | | |
| Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District | | |
| Wuxi Jiangsu Province, China | | |
| TELE: 86-510-82732223 FAX: 86-510-82720101-8014 | | |
| Web : <u>www.dldevelop.com</u> | | |
| Email: info@dldevelop.com.cn service@dldevelop.com.cn | | |
| | | |
| Anti-Human Ig | light chain λ | Monoclonal |
| Antibody(PE Conjugated) | | |
| DL21902F | | |
| | | |
| Immunoglobulin light chain lambda,IGL | | |
| U | 0 | , |
| The MHL-38 antibody reacts with both soluble and | | |
| membrane human immunoglobulin light chain | | |
| lambda (λ). It does not react with human | | |
| immunoglobulin light chain kappa (κ) or heavy | | |
| chains. The MHL-38 antibody can be used as | | |
| primary or secondary reagent for immunofluorescent | | |
| | | |
| 8 | | |
| | Isotype: N | Mouse IgG2a, к |
| | ~ 1 | Č , |
| Size: 20 Tests/50 Tests/100 Host: Mouse | | |
| | | |
| | V SCI&TECH DI Mingpin CityII,I vince, China 2732223 relop.com velop.com.cn Anti-Human Ig Antibody(PE Co DL21902F Immunoglobulin The MHL-38 ar membrane huma lambda (λ). It do immunoglobulin chains. The MH primary or secon staining or ELIS | V SCI&TECH DEVELOPME Mingpin CityII,No.8 Xihudo vince, China 2732223 FAX: relop.com velop.com.cn service@dld Anti-Human Ig light chain λ Antibody(PE Conjugated) DL21902F Immunoglobulin light chain The MHL-38 antibody reacts membrane human immunogl lambda (λ). It does not react immunoglobulin light chain chains. The MHL-38 antibod primary or secondary reagen staining or ELISA analysis. Isotype: I |



Tests/200 Tests Reactivity: Human

Application: FCM

Concentration:

Conjugation: PE

SwissProt:

Buffer: PBS with 0.05% Proclin300, 1% BSA

Recommended Use:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data:

Human peripheral blood lymphocytes are stained with Anti-Human Ig light chain λ Monoclonal Antibody(PE Conjugated)(filled gray histogram). Unstained lymphocytes (empty black histogram) are used as control.

Storage:

Keep as concentrated solution.



Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.

