

Manufacturer/Supplier:		
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD		
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District		
Wuxi Jiangsu Province, China		
TELE: 86-510-8	2732223	FAX: 86-510-82720101-8014
Web : <u>www.dldevelop.com</u>		
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>		
Product name:	Anti-Human Igl	M Monoclonal Antibody(FITC
	Conjugated)	
Catalog:	DL21914F	
Synonyms:	Immunoglobulin heavy constant mu,IGHM,Immunoglobulin M	
Background:	ckground: IgM is the first immunoglobulin made by B cells in the immune response. Surface IgM is expressed on immature and mature B cells, while IgM heavy (μ) chain is expressed intracellularly in pre-B cells.	
Form: Liquid		Isotype: Mouse IgG1, κ
-		
Size:		Host: Mouse
20Tests/100Tests/100Tests×2		
Reactivity: Human		Application: FCM
v		~ *



Concentration: 5 µL

Conjugation: FITC

SwissProt: P01871

Buffer:Phosphate buffered solution, pH 7.2, containing
0.09% stabilizer and 1% protein protectant.

Recommended Use:


Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data:

Human peripheral blood lymphocytes are stained with Anti-Human IgM Monoclonal Antibody(FITC Conjugated)(filled gray histogram) or Mouse IgG1 Isotype Control FITC (empty black histogram).



DEVELOP



Storage:

Keep as concentrated solution.

Store at $2\sim 8^{\circ}$ C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.

