

**Manufacturer/Supplier:**

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

**Address:** A1-203 Mingpin CityII, No.8 Xihudong Road, Liangxi District  
Wuxi Jiangsu Province, China

TELE: 86-510-82732223

FAX: 86-510-82720101-8014

**Web :** [www.dldevelop.com](http://www.dldevelop.com)

Email: [info@dldevelop.com.cn](mailto:info@dldevelop.com.cn)    [service@dldevelop.com.cn](mailto:service@dldevelop.com.cn)

**Product name:** Anti-Human IL-17A Monoclonal Antibody(FITC  
Conjugated)

**Catalog:** DL21923F

**Synonyms:** CTLA8, IL17, CTLA-8, IL 17A

**Background:** IL-17A is the founding member of the IL-17 family, a group of six structurally related pro- inflammatory cytokines. IL-17A, secreted by activated CD4<sup>+</sup> Th17 cell subpopulation, elicits multiple biological activities on a variety of cells including: the induction of IL-6, IL-8, G-CSF, and PGE2 production in epithelial, endothelial or fibroblasts; the enhancement of surface expression of ICAM-1 in fibroblasts; activation of NF- $\kappa$ B and costimulation of T cell proliferation. Recent studies demonstrated that, in mice, activated IL-17-secreting CD4<sup>+</sup> helper T

cells (Th17 cells) mediate an autoimmune arthritis that clinically and immunologically resembles rheumatoid arthritis (RA). Human IL-17A shows 63%, 63%, and 72% amino acid sequence identity to rat IL-17A, mouse IL-17A, and a protein encoded by the ORF13 gene of herpesvirus Saimiri (HVS), respectively.

**Form:** Liquid

**Isotype:** Mouse IgG1,  $\kappa$

**Size:**

**Host:** Mouse

20Tests/100Tests/100Tests $\times$ 2

**Reactivity:** Human

**Application:** ICFCM

**Concentration:** 5  $\mu$ L

**Conjugation:** FITC

**SwissProt:** Q16552

**Buffer:** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

### Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per

100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

**Storage:**

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.