

| Manufacturar/Si | unlier | | |
|---|---|--|---|
| Manufacturer/Supplier: | | | |
| WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD | | | |
| Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District | | | |
| Wuxi Jiangsu Province, China | | | |
| TELE: 86-510-82732223 FAX: 86-510-827201 | | | |
| Web : <u>www.dldevelop.com</u> | | | |
| Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u> | | | |
| | | | |
| Product name: | Anti-Mouse CD31 Monoclonal | | |
| | Antibody(PerCP/Cyanine5.5 Conjugated) | | |
| Catalog: | DL21991F | | |
| | | | |
| Synonyms: | Pecam, Pecam-1, PECAM-1, CD31 | | |
| | | | |
| Background: | CD31 is a 130-140 kD glycoprotein, also known as | | |
| | platelet endothelial cell adhesion molecule (PECAM-1) and EndoCAM. It is a member of the Ig superfamily, expressed on endothelial cells, platelets, | | |
| | | | |
| | | | |
| granulocytes, monocytes/macrophages, dendritic cells, and T and B cell subsets, and is critical for | | | |
| | | | cell-cell interactions. The primary ligands for CD3 |
| | have been reported to be CD38 and the vitronectin | | |
| | receptor ($\alpha v \beta 3$ integrin, CD51/CD61). Other | | |
| | | | |
| | reported functions of CD31 are neutrophil emigration | | |
| | to sites of inflammation and angiogenesis. | | |



| Form: Liquid | | Isotype: Rat IgG2a, κ |
|--------------------------|---|-------------------------------|
| Size: 25µg/100µg | | Host: Rat |
| Reactivity: Mouse | | Application: FCM |
| Concentration: 0.2 mg/mL | | Conjugation: PerCP/Cyanine5.5 |
| SwissProt: | Q08481 | |
| Buffer: | Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant. | |

Recommended Use:


Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 μ g/10⁶cells in 100 μ L volume].

Data:

C57BL/6 murine splenocytes are stained with Anti-Mouse CD31 Monoclonal Antibody(PerCP/Cyanine5.5 Conjugated)[Used at 0.2 μ g/10⁶ cells dilution](filled gray histogram). Unstained splenocytes



(empty black histogram) are used as control.



Storage:

Keep as concentrated solution.

Store at $2\sim 8^{\circ}$ C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.