

Manufacturer/Supplier:			
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD			
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District			
Wuxi Jiangsu Province, China			
TELE: 86-510-82732223 FAX: 86-510-82720101-8014			86-510-82720101-8014
Web : <u>www.dldevelop.com</u>			
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>			
Product name:	U	Monoclona	l Antibody(FITC
Catalog:	Conjugated) DL22067F		
Synonyms:	Immunoglobulin heavy constant delta,IGHD,Ig delta chain C region		
Background:	Surface IgD is an important B cell differentiation marker.		
Form: Liquid		Isotype:	Rat IgG2a, к
Size: 25µg/100µg		Host: Rat	
Reactivity: Mouse		Application: FCM	
Concentration: 0.5 mg/mL		Conjugat	ion: FITC

DEVELOP

SwissProt: P01881

Buffer:Phosphate buffered solution, pH 7.2, containing
0.09% stabilizer and 1% protein protectant.

Recommended Use:


Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 μ g/10⁶cells in 100 μ L volume].

Data:

C57BL/6 murine splenocytes are stained with Anti-Mouse IgD Monoclonal Antibody(FITC Conjugated)[Used at $0.2 \ \mu g/10^6$ cells dilution](filled gray histogram). Unstained splenocytes (empty black histogram) are used as control.



DEVELOP



Storage:

Keep as concentrated solution.

Store at $2\sim 8^{\circ}$ C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.