

Manufacturer/Supplier:			
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD			
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District			
Wuxi Jiangsu Province, China			
TELE: 86-510-82732223 FAX: 86-510-82720101-8014			
Web : <u>www.dldevelop.com</u>			
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>			
Product name:	Anti-Mouse Ly6A/E(Sca-1) Monoclonal		
	Antibody(FITC Conjugated)		
Catalog:	DL22094F		
Synonyms:	Ly6a,Ly6,Ly-6A.2/Ly-6E.1,SCA-1,TAP		
Background:	Ly-6A/E, also known as Sca-1, is an 18 kD member		
	of the Ly-6 multigene family. Ly6A/E is a		
	glycosylphosphatidylinositol (GPI)-linked protein		
	expressed on hematopoietic stem cells. In mice		
	expressing the Ly-6.2 haplotype (e.g., AKR, C57BL,		
	C57BR, DBA/2, SJL, SWR, and 129), Ly-6A/E is		
	also expressed on peripheral B lymphocytes and		
	thymic and peripheral T lymphocytes. Strains		
	expressing the Ly-6.1 haplotype (e.g., BALB/c,		
	CBA, C3H/He, DBA/1, and NZB) have low Ly-6A/E		
	expression on resting peripheral lymphocytes. The		



expression of Ly-6A/E on lymphocytes is upregulated upon activation from both Ly6.1 and Ly6.2 haplotype mice. Ly-6A/E is thought to be involved in the regulation of both T and B cell responses.

Form: Liquid		Isotype: Rat IgG2a, κ
Size: 50Tests/100Tests/100Tests×2		Host: Rat
Reactivity: Mouse		Application: FCM
Concentration: 5 µL		Conjugation: FITC
SwissProt:	P05533	
Buffer:	Phosphate buffered solution, pH 7.2, containing	

Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined


0.09% stabilizer and 1% protein protectant.

DEVELOP

for individual use.

Data:

C57BL/6 murine splenocytes are stained with Anti-Mouse Ly6A/E(Sca-1) Monoclonal Antibody(FITC Conjugated)(filled gray histogram). Unstained splenocytes (empty black histogram) are used as control.



Storage:

Keep as concentrated solution.

Store at $2\sim 8^{\circ}$ C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.

