

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Mouse CD38 Monoclonal Antibody(FITC

Conjugated)

Catalog: DL22120F

Synonyms: ADP-ribosyl cyclase 1,ADPRC 1,CD38,NIM-R5

antigen, 2'-phospho-cyclic-ADP-ribose transferase

Background: CD38 is a 42 kD glycoprotein, also known as T10. It

> is an ADP-ribosyl hydrolase, expressed on B cells, NK cells, a subset of T cells, brain, muscle, and

kidney. In mouse, CD38 expression is downregulated on germinal center B cells and plasma cells, whereas

this is not the case for humans. By functioning as both a cyclase and a hydrolase, CD38 mediates lymphocyte activation, as well as adhesion and

metabolism of cADPR and NAADP. CD31 is the

ligand of CD38.



Form: Liquid **Isotype:** Rat IgG2a, κ

Size: Host: Rat

50Tests/100Tests/100Tests×2

Reactivity: Mouse **Application:** FCM

Concentration: 5 µL **Conjugation: FITC**

SwissProt: P56528

Phosphate buffered solution, pH 7.2, containing Buffer:

0.09% stabilizer and 1% protein protectant.

Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Storage:

Keep as concentrated solution.



Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.