

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road, Liangxi District
Wuxi Jiangsu Province, China

TELE: 86-510-82732223

FAX: 86-510-82720101-8014

Web : www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Mouse IL-10 Monoclonal Antibody(APC
Conjugated)

Catalog: DL22156F

Synonyms: Interleukin-10, IL-10, Cytokine synthesis inhibitory
factor, CSIF

Background: L-10 was originally described as Cytokine Synthesis
Inhibitory Factor (CSIF) by virtue of its ability to
inhibit cytokine production by Th1 clones. IL-10
shares over 80% sequence homology with the
Epstein-Barr virus protein BCRF1. IL-10 inhibits
IFN- γ , TNF- β , and IL-2 production by Th1 clones;
inhibits macrophage-mediated IL-1, IL-6, and TNF- α
synthesis; suppresses the delayed type
hypersensitivity response; stimulates Th2 cell
response (which results in elevated antibody

production); and promotes mast cell proliferation in combination with IL-4.

Form: Liquid

Isotype: Rat IgG2b, κ

Size: 25 μ g/100 μ g

Host: Rat

Reactivity: Mouse

Application: ICFCM

Concentration: 0.2 mg/mL

Conjugation: APC

SwissProt: P18893

Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 μ g/10⁶cells in 100 μ L volume].

Storage:

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.