

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road, Liangxi District
Wuxi Jiangsu Province, China

TELE: 86-510-82732223

FAX: 86-510-82720101-8014

Web : www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Mouse H-2 Monoclonal Antibody(APC
Conjugated)

Catalog: DL22271F

Synonyms: Mouse major histocompatibility complex (MHC)
H-2, MHC I

Background: The M1/42 antibody reacts with the H-2 MHC class I
alloantigens expressed on nucleated cells from mice
of the a, b, d, j, k, s, and u haplotypes (Stallcup, KC
et al, 1981). MHC class I is involved in antigen
presentation to T cells expressing CD3/TCR and
CD8 proteins.

Form: Liquid

Isotype: Rat IgG2a, κ

Size:

Host: Rat

50Tests/100Tests/100Tests×2

Reactivity: Mouse

Application: FCM

Concentration: 5 µL

Conjugation: APC

SwissProt:


Buffer: Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 µL staining volume or per 100 µL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data:

C57BL/6 murine splenocytes are stained with Anti-Mouse H-2 Monoclonal Antibody(APC Conjugated)(filled gray histogram). Unstained splenocytes (empty black histogram) are used as control.



Storage:

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.