

## Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

**Product name:** Anti-Mouse H-2 Monoclonal Antibody(APC

Conjugated)

Catalog: DL22280F

**Synonyms:** Mouse major histocompatibility complex (MHC)

H-2, MHC I

**Background:** The M1/42 antibody reacts with the H-2 MHC class I

alloantigens expressed on nucleated cells from mice of the a, b, d, j, k, s, and u haplotypes (Stallcup, KC et al, 1981). MHC class I is involved in antigen presentation to T cells expressing CD3/TCR and

CD8 proteins.

Form: Liquid Isotype: Rat IgG2a, κ

Size: 25µg/100µg Host: Rat



**Reactivity:** Mouse **Application:** FCM

**Concentration:** 0.2 mg/mL **Conjugation:** APC

SwissProt:

Phosphate buffered solution, pH 7.2, containing Buffer:

0.09% stabilizer and 1% protein protectant.


## Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1  $\mu g/10^6$  cells in 100  $\mu L$  volume].

## Data:

C57BL/6 murine splenocytes are stained with Anti-Mouse H-2 Monoclonal Antibody(APC Conjugated)[Used at 0.2 µg/10<sup>6</sup> cells dilution](filled gray histogram). Unstained splenocytes (empty black histogram) are used as control.





## **Storage:**

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.